

FRONT / CLOSE

FRONT / CLOSE

95mm (w) x 60mm(h)

BACK / CLOSE

FOR HEALTHCARE PROFESSIONALS

IMPORTANT SAFETY INFORMATION YOU SHOULD KNOW:
TECVAYLI® and **TALVEY®** therapy can cause cytokine release syndrome (CRS) or neurologic toxicity, including Immune Effector Cell-Associated Neurotoxicity Syndrome (ICANS) which may be fatal or life-threatening. CRS may involve multiple organ systems.



TECVAYLI® Prescribing
Information



TALVEY® Prescribing
Information

Complete the Patient Card based on the drug prescribed and provide the Patient Card to the patient.

This Patient has received: (Please check one) ☐ **TECVAYLI®** (teclistamab) ☐ **TALVEY®** (talquetamab)

Name of Treating Physician: _____

Treating Physician's Clinic Contact Number: _____

Hospital Name: _____

Hospital Phone Number: _____

Dosing Schedule: (Please check one) ☐ **Weekly Dosing** ☐ **Bi-Weekly Dosing**

Dates of Infusions:

Step-up Dose 1: _____ **Step-up Dose 2:** _____

Step-up Dose 3: _____ **First Treatment Dose:** _____

*Applies only if patient is receiving TALVEY® at a treatment dose of 0.8 mg/kg
 For more information about TECVAYLI® or TALVEY®, please scan the QR code for the Singapore Package Insert
 Tecvayli & Talvey Patient Card Version 1. This document has been approved by HSA as of 15-Nov-2024

EM-171289

INSIDE / OPEN



FOR THE PATIENT

Call your healthcare professional or get emergency help right away if you recognize any of these symptoms:

Symptoms of Cytokine Release Syndrome (CRS):	Symptom of neurologic toxicity including ICANS:
- Fever (38°C or higher) - Difficulty breathing - Chills - Fast heartbeat - Dizziness or light-headedness - Confusion or restlessness - Headache - Low blood pressure	- Headache - Feeling confused or disoriented - Being less alert or aware - Feeling sleepy - Low energy (lethargy) - Slowness or difficulty thinking - Seizures - Memory Loss - Jerking movements - Rigid muscles - Feeling restless - Numbness and tingling (feeling like "pins and needles") - Trouble speaking or writing - Muscle spasms - Shaking (tremors) - Problems walking - Hearing Loss - Muscle weakness in your body or face - Double vision - Burning, throbbing or stabbing pain

You should always ask your doctor about taking other medications while taking **TECVAYLI®** or **TALVEY®**.



IMPORTANT TO REMEMBER: Stay close to the location where you received your therapy for at least 2 days for daily monitoring after administration of all doses within the step-up dosing schedule. If you have any of these symptoms, call your doctor or seek emergency medical attention right away! These are not all of the possible symptoms of **TECVAYLI®** or **TALVEY®**. Tell your doctor if you have any symptom that bothers you or does not go away.

Johnson & Johnson

Johnson & Johnson International (Singapore) Pte Ltd
No 2, Science Park Drive, #07-13, Ascent, Science Park 1, Singapore 118222



(teclistamab)



(talquetamab)

95mm (w) x 119mm(h)